arXiv:1612.03509v1 [hep-ph] 12 Dec 2016

# The energy loss effect of incoming quark in the Drell-Yan reaction off nuclei

Li-Hua Song,[1,*] Lin-Wan Yan,[1] and Chun-Gui Duan[2,†]

[1] *College of Science, North China University of Science and Technology, Tangshan 063009, P.R.China*

[2] *Department of Physics, Hebei Normal University, Shijiazhuang 050024, P.R.China*

## Abstract

By means of the SW quenching weights and the analytic parameterizations of quenching weights based on BDMPS formalism, the leading-order computations for nuclear Drell-Yan differential cross section rations as a function of the quark momentum fraction are performed with the nuclear geometry effect in Drell-Yan dimuon production. The transport coefficient of the incoming quark energy loss in cold nuclear matter is extracted with a global analysis of the experimental data from NA3 and E866 collaborations. The extracted transport coefficient with the SW quenching weights is $\hat{q} = 0.32 \pm 0.04$ GeV$^2$/fm, which is approximately equal to the value $\hat{q} = 0.37 \pm 0.05$ GeV$^2$/fm determined with the analytic parameterizations of BDMPS quenching weights. It is found that our predictions are in good agrement with the experimental measurements, and especially the agreement of calculations with NA3 experimental data has a significant improvement. The comparison of the theoretical results and the corresponding experimental data demonstrates intuitively that the energy loss effect of incoming quark is particularly significant at lower incident proton energies. It is hoped that the obtained results relevant to the NA3 experiment ($E_{beam} = 150$ GeV) can provide useful reference for the forthcoming E906 Sea Quest experiment ($E_{beam} = 120$ GeV).

**Keywords:** energy loss, quark distribution, Drell-Yan process

PACS numbers: 12.38.-t;13.85.Qk;24.85.+p;25.40.-h

*E-mail: songlh@ncst.edu.cn

†E-mail: duancg@mail.hebtu.edu.cn

## I. INTRODUCTION

A series of experimental studies on jet-quenching from RHIC[1-6] and LHC[7] reflect the energy loss of fast partons while traversing the hot and dense medium. The parton energy loss in high energy collisions has attracted attention from both the nuclear and particle physics communities during the last two decades. However, any consensus in the baseline for the parton energy loss has been lacking. A solid understanding of the nuclear modification of particle spectra in cold nuclear matter is fundamentally important for investigating the fast parton propagation in hot and dense medium and further quantifying the properties of the QGP created in these heavy-ion collisions.

The Drell-Yan reaction[8] in hadron-nucleus collisions can provide the possibility of probing the parton propagation through nuclear matter, with the produced lepton pair carrying away the desired information on the parton after it has travelled in the nucleus. As the lepton pair in the final state does not interact strongly with partons in the nuclei, only initial-state interactions are important in the Drell-Yan scattering. The Drell-Yan process can be considered as a very clean probe for the energy loss effect of incoming quark in cold nuclear matter. A series of experimental data on the Drell-Yan differential cross section ratios were presented by NA3[9] and NA10[10] Collaborations from CERN, and E772[11] and E866[12] Collaborations from Fermilab. The data on the differential cross section ratio as a function of the quark momentum fraction can cancel the most uncertainties regarding the lepton pair production, and avoid the influence of the QCD next-to-leading order correction[13]. Many theoretical phenomenological models were proposed to describe the incoming quark energy loss in nuclear Drell-Yan process[14-19]. The best value of energy loss obtained by their independent analysis of the Drell-Yan experimental data was strongly dependent on the used nuclear parton distribution functions and the assumption for the average path length of the incident quark.

In our preceding papers[20-23], the effect of energy loss in Drell-Yan process is discussed and models are given to incorporate them in the calculation of Drell-Yan yields. It is found that the energy loss effect of incoming quark can suppress evidently the differential cross sections versus the quark momentum fraction, and the mean value of quark energy loss obtained by HKM nuclear parton distribution functions[24] is larger than that by using HKN07[25], nDS[26] and EPS09[27] parameterizations which directly reflects the deviation

between HKM nuclear corrects to sea quark distribution and other sets. It is noticeable that HKM proposed the nuclear parton distributions that were determined only by fitting the experimental data from $l$-A deep inelastic scattering (DIS) without the data about Drell-Yan reaction. It is worth to mention that the mean energy loss employed in our preceding calculations wold be considered very simplistic in hot and dense matter, and the fluctuations of the path that a quark takes through the medium has not yet been considered.

It is accepted that at least the probability $D(\varepsilon, L)$ of energy loss $\varepsilon$ given a path $L$ with averaged over geometry is relevant quantity. In this paper, by means of the Salgado-Wiedemann (SW) quenching weights[28], and the analytic parameterizations of quenching weights[29] based on Baier, et al. (BDMPS) formalism[30], the leading-order computations for nuclear Drell-Yan differential cross section ration as a function of the quark momentum fraction are performed with the nuclear geometry effect in Drell-Yan dimuon production. The transport coefficient $\hat{q}$ of the incoming quark energy loss in cold nuclear matter is extracted with a global analysis of the NA3[9] (incident proton energy $E_{beam} = 150$ GeV and target-quark momentum fraction $x_2$ from 0.074 to 0.366) and E866[12] ($E_{beam} = 800$ GeV and $0.0179 < x_2 < 0.1022$) nuclear Drell-Yan experimental data. We desire that our results and discussion in this work can facilitate the theoretical research on the microscopic dynamics of medium-induced parton energy loss.

The remainder of this paper is organized as follows. In Section II, the theoretical framework of our study is introduced. Section III is devoted to the results and discussion. Finally, a summary is presented.

## II. DRELL-YAN REACTION IN HADRON-NUCLEUS COLLISIONS

At the leading order in perturbation theory, the lepton pair production differential cross section in hadron-nucleus collisions is obtained from the convolution of differential partonic cross section $q\bar{q} \to l^+l^-$ with the quark distributions in the beam and in the target:

$$\frac{d^2\sigma}{dx_1dx_2} = \frac{4\pi\alpha_{em}^2}{9sx_1x_2}\sum_f e_f^2[q_f^h(x_1,Q^2)\bar{q}_f^A(x_2,Q^2) + \bar{q}_f^h(x_1,Q^2)q_f^A(x_2,Q^2)]. \tag{1}$$

where $x_1(x_2)$ is the momentum fraction carried by the projectile (respectively target) parton, $\alpha_{em}$ is the fine structure constant, $\sqrt{s}$ is the center of mass energy of the hadronic collision, $e_f$ is the charge of the quark with flavor $f$, $Q^2$ is the invariant mass of a lepton pair, the

sum is carried out over the light flavor ($f = u, d, s$), and $q_f^{h(A)}(x, Q^2)$ and $\bar{q}_f^{h(A)}(x, Q^2)$ are respectively the quark and anti-quark distribution function in the hadron (nucleon in the nucleus A).

In the hadron-induced Drell-Yan reaction on nucleus, when the incoming quark propagates through the nuclear medium, it suffers multiple scattering on the surrounding nucleon and gluon radiation. The induced gluon bremsstrahlung effectively reduces the incoming quark energy. The energy loss ($\varepsilon$) of the incoming quark results in a change in its momentum fraction prior to the collision, $\Delta x_1 = \varepsilon / E_{beam}$, where $E_{beam}$ is the incident hadron energy. As a consequence, the projectile parton distribution function should be evaluated at $x_1' = x_1 + \Delta x_1$.

With considering the incoming quark energy loss in the nuclear medium, the Drell-Yan differential cross section in hadron-nucleus collisions can be expressed as:

$$\frac{d^2\sigma}{dx_1 dx_2} = \frac{4\pi\alpha_{em}^2}{9 s x_1 x_2} \sum_f e_f^2 \int_0^{\varepsilon_{max}} d\varepsilon D(\varepsilon, \omega_c, L)[q_f^h(x_1', Q^2)\bar{q}_f^A(x_2, Q^2) + \bar{q}_f^h(x_1', Q^2) q_f^A(x_2, Q^2)] \quad (2)$$

Here the characteristic gluon frequency $\omega_c = \frac{1}{2}\hat{q}L^2$, and $D(\varepsilon, \omega_c, L)$ denotes the probability that an incoming quark suffers an energy loss $\varepsilon$ from the quark radiating gluons. In the calculation, the energy scale $\omega_c$ depends on the path length $L$ covered by the incoming quark in the nuclear medium, and is determined by the transport coefficient $\hat{q}$ adjusted to the data.

Considering the fluctuations of the path $L$ travelled by the incoming quark across the target, the energy loss probability distribution $D(\varepsilon, \omega_c, L)$ needs to be averaged over geometry. The nuclear geometry is described by the nuclear density distribution $\rho_A(\vec{b}, y)$, where $y$ is the coordinate along the direction of the incoming quark and $\vec{b}$ is the impact parameter. The center of the target nucleus lies at $(\vec{0}, 0)$. With the assumption that the annihilation of a quark-antiquark pair into a virtual photon is located at $(\vec{b}, y)$, the incident quark at $y$ will travel the path length $L = \sqrt{R_A^2 - b^2} + y$, along a direction with impact parameter $\vec{b}$. Then with the nuclear geometry effect in Drell-Yan dimuon production, the nuclear Drell-Yan differential cross section should be written as:

$$\frac{d^2\sigma^{h-A}}{dx_1 dx_2} = \frac{4\pi\alpha_{em}^2}{9 s x_1 x_2} \sum_f e_f^2 \int d^2b dy \rho_A(\vec{b}, y) \int_0^{\varepsilon_{max}} d\varepsilon D(\varepsilon, \omega_c, L) \times [q_f^h(x_1', Q^2)\bar{q}_f^A(x_2, Q^2) + \bar{q}_f^h(x_1', Q^2) q_f^A(x_2, Q^2)]. \quad (3)$$

Here the nuclear density profile $\rho_A(\vec{b}, y)$ is normalized to unity.

## III. RESULTS AND DISCUSSION

In order to extract the value of the transport coefficient $\hat{q}$, we give a phenomenological analysis at leading order for the nuclear Drell-Yan differential cross section ratio:

$$R_{A_1/A_2}(x_{1(2)}) = \int dx_{2(1)} \frac{d^2\sigma^{h-A_1}}{dx_1 dx_2} / \int dx_{2(1)} \frac{d^2\sigma^{h-A_2}}{dx_1 dx_2}, \tag{4}$$

for the NA3[9] and E866[12] experimental data (87 points). By means of the CERN subroutine MINUIT[31], the transport coefficient $\hat{q}$ is obtained by minimizing $\chi^2$. One standard deviation of the optimum parameter correspond to an increase of $\chi^2$ by 1 unit from its minimum $\chi^2_{min}$. In our calculation, we use the uniform hard-sphere nuclear density normalized to unity,

$$\rho_A(\sqrt{b^2+y^2}) = (\rho_0/A)\Theta(R_A - \sqrt{b^2+y^2}). \tag{5}$$

Here $\rho_0$ is the nuclear density, $R_A = r_0 A^{1/3}$, and $r_0 = 1.12$ fm.

Table I summarizes the calculated results corresponding to the transport coefficient $\hat{q}$ and $\chi^2$ per number of degrees of freedom ($\chi^2/ndf$) by means of the SW quenching weights[28] and the HKM nuclear parton distribution functions[24]. It is shown that the global fit of all data gives $\hat{q} = 0.32 \pm 0.04$ GeV$^2$/fm with $\chi^2/ndf = 0.90$. As can be seen from Table 1, considering the incoming quark energy loss effect and the nuclear geometry effect in Drell-Yan dimuon production, the theoretical results are in good agreement with the experimental data, and especially the agreement of calculations with NA3[9] experimental data has a significant improvement (with $\chi^2/ndf = 0.54$ for NA3$x_1$ data and $\chi^2/ndf = 1.28$ for NA3$x_2$ data). In our previous study [23], $\chi^2/ndf$ extracted from the NA3$x_{1(2)}$ data is 1.69 (respectively 2.53) by means of the mean energy loss model and without the fluctuations of the path.

To demonstrate intuitively the energy loss effect of an incoming quark on the nuclear Drell-Yan cross section ratio, the solid curves in Fig.1, Fig.2, Fig.3 and Fig.4 are our numeric results which are compared with the corresponding experimental data. In Fig.1, it is necessary to note that NA3 Collaboration provided the negative pion incident Drell-Yan cross section ratio $R_{H/Pt}$ as a function of parton momentum fraction. It is shown that the theoretical results with the incoming quark energy loss effect and the nuclear geometry effect in Drell-Yan dimuon production have good agreement with the experimental data. As can be seen from Fig.1 that the influence due to the incoming quark energy loss on the NA3

TABLE I: The values of $\hat{q}$ and $\chi^2/ndf$ extracted from the experimental data by means of the SW quenching weights[28].

| Exp.data | $\hat{q}$ (GeV$^2$/fm) | $\chi^2/ndf$ |
|---|---|---|
| E866$x_1$ | $0.29 \pm 0.05$ | 0.86 |
| E866$x_2$ | $0.41 \pm 0.08$ | 1.18 |
| NA3$x_1$ | $0.42 \pm 0.11$ | 0.54 |
| NA3$x_2$ | $0.50 \pm 0.09$ | 1.28 |
| Global fit | $0.32 \pm 0.04$ | 0.90 |

nuclear Drell-Yan different cross section ratio ($R_{H/Pt}(x_1)$) decreases with the increase of $x_1$ (approximately from 69% to 24% in the range $0.25 < x_1 < 0.65$) and then becomes greater (approximately from 24% to 68% in the range $0.65 < x_1 < 0.95$). As for the ratio $R_{H/Pt}(x_2)$, the influence induced by the quark energy loss is also firstly decreases with the increase of $x_2$ (approximately from 54% to 33% in the range $0.07 < x_2 < 0.17$), and then increase approximately from 33% to 49% in the range $0.17 < x_2 < 0.37$. From Fig.2, Fig.3 and Fig.4, it is found that the incoming quark energy loss effect on the Fermilab E866 nuclear Drell-Yan different cross section ratio becomes greater with the increase of quark momentum fraction ($x_1$) in the beam hadron (for example, the suppression of $R_{W/Be}(x_1)$ for $M = 4 - 5$ is approximately from 4% to 19% in the range $0.25 < x_1 < 0.95$ ), and decreases gradually with the increase of quark momentum fraction ($x_2$) in the target nucleus (such as the suppression of $R_{W/Be}(x_2)$ is approximately from 10% to 3% in the range $0.018 < x_2 < 0.102$ ). It is obvious that the energy loss effect of incoming quark is particularly significant at lower incident proton energies. It is worth emphasizing that shadowing should also occur in nuclear Derll-Yan process at small $x_2$ ($x_2 < 0.15$). High statistics , high precision data on nuclear Drell-Yan different cross section ratio from lower incident proton energies in larger $x_2$ region will help to pin down the energy loss effect of an incoming quark in cold nuclear matter.

In Table II, we presents the results obtained by means of the analytic parameterizations of BDMPS quenching weights[29]:

$$D(\varepsilon, \omega_c, L) = \frac{1}{\sqrt{2\pi}\sigma(\varepsilon/\omega_c)} \exp[-\frac{(\log(\varepsilon/\omega_c) - \mu)^2}{2\sigma^2}], \tag{6}$$




FIG. 1: The nuclear Drell-Yan cross section ratios $R_{H/Pt}(x_{1(2)})$ calculated with SW quenching weights[28]. The experimental data are taken from the NA3[9].




FIG. 2: The nuclear Drell-Yan cross section ratios $R_{W/Be}(x_1)$ calculated with SW quenching weights[28]. The experimental data are taken from the E866[12]

where $\mu = -2.55$ and $\sigma = 0.57$. As can be seen from Table II, the global fit of all data gives $\hat{q} = 0.37 \pm 0.05$ GeV$^2$/fm with $\chi^2/ndf = 0.96$, which is approximately equal to the above result ($\hat{q} = 0.32 \pm 0.04$ GeV$^2$/fm with $\chi^2/ndf = 0.90$ ) given by the SW quenching weights[28]. By comparing with our previous discussion on the semi-inclusive deep inelastic scattering of




FIG. 3: The nuclear Drell-Yan cross section ratios $R_{Fe/Be}(x_1)$ calculated with SW quenching weights[28]. The experimental data are taken from the E866[12]




FIG. 4: The nuclear Drell-Yan cross section ratios $R_{A/Be}(x_2)$ calculated with SW quenching weights[28]. The experimental data are taken from the E866[12]

lepton on nuclear targets[32], the transport coefficient ($\hat{q} = 0.20 \pm 0.02$ GeV$^2$/fm) extracted from the selected HERMES experimental data with the quarks hadronization occurring outside the nucleus by means of the analytic parameterizations of BDMPS quenching weights is much smaller than that obtained by the SW quenching weights ($\hat{q} = 0.74 \pm 0.03$ GeV$^2$/fm).

TABLE II: The values of $\hat{q}$ and $\chi^2/ndf$ extracted from the experimental data by means of the analytic parameterizations of BDMPS quenching weights[29].

| Exp.data | $\hat{q}$ (GeV$^2$/fm) | $\chi^2/ndf$ |
|---|---|---|
| E866$x_1$ | $0.37 \pm 0.05$ | 0.92 |
| E866$x_2$ | $0.36 \pm 0.12$ | 1.08 |
| NA3$x_1$ | $0.40 \pm 0.07$ | 1.49 |
| NA3$x_2$ | $0.41 \pm 0.09$ | 1.08 |
| Global fit | $0.37 \pm 0.05$ | 0.96 |

As discussed in paper[32], the finite-length effects which lie in the SW quenching weights result in the large value of the transport coefficient $\hat{q}$ for small-$R_A$ target. This indicates that the finite-length effects have little influence on extracting the transport coefficient $\hat{q}$ from E866 and NA3 Drell-Yan experimental data due to the large-$R_A$ target. In addition, the value of the transport coefficient ($\hat{q} = 0.37 \pm 0.05$ GeV$^2$/fm) obtained in this work for incoming quark by means of the analytic parameterizations of BDMPS quenching weights is bigger than that ($\hat{q} = 0.20 \pm 0.02$ GeV$^2$/fm) extracted from the HERMES experimental data for outgoing quark (with $\mu = -1.5$ and $\sigma = 0.73$ in eq.6). Here, the uncertainties of the HKM nuclear parton distributions can be the main source of uncertainties associated with our calculated results. The precise measurements from E906 Sea Quest[33] experiments on nuclear Drell-Yan different cross section ratio in various regions of $x_1$ and $x_2$ will provide important information about the modification of quark PDFs and nucleon structure function in the nucleus, and further constrain the transport coefficient in cold nuclear medium.

## IV. SUMMARY

The investigation into the fast parton energy loss in cold nuclear matter is crucial for a solid understanding of the parton propagation in hot-dense medium. The Drell-Yan reaction in hadron-nucleus is an ideal tool to determine and constrain the transport coefficient in cold nuclear medium. By means of the SW quenching weights and the analytic parameterizations of quenching weights based on BDMPS formalism, the NA3 and E866 experimental data on nuclear Drell-Yan differential cross section ratio as a function of the quark momentum

fraction have been analyzed with the HKM nuclear parton distribution functions. Considering the fluctuations of the path $L$ travelled by the incoming quark across the target, the energy loss probability distribution $D(\varepsilon, \omega_c, L)$ has been averaged over geometry. Due to the large-$R_A$ target, the finite-length effects which lie in the SW quenching weights have little influence on extracting the transport coefficient $\hat{q}$ from NA3 and E866 nuclear Drell-Yan experimental data. The extracted transport coefficient from the global fit is shown to be $\hat{q} = 0.32 \pm 0.04$ GeV$^2$/fm ($\chi^2/ndf = 0.90$ ) for the SW quenching weights, which is approximately equal to the value $\hat{q} = 0.37 \pm 0.05$ GeV$^2$/fm ($\chi^2/ndf = 0.96$) determined with the analytic parameterizations of BDMPS quenching weights. It is found that our predictions are in good agrement with the experimental measurements, and especially the agreement of calculations with NA3 experimental data has a significant improvement. The comparing with the corresponding experimental data demonstrates intuitively that the energy loss effect of incoming quark is particularly significant at lower incident proton energies. It is worth emphasizing that shadowing should also occur in nuclear Derll-Yan process at small $x_2$ ($x_2 < 0.15$). The uncertainties of the HKM nuclear parton distributions can be the main source of uncertainties associated with our calculated results. High statistics , high precision data from E906 Sea Quest experiment on nuclear Drell-Yan different cross section ratio at $E_{beam} = 120$ GeV in various regions of $x_1$ and $x_2$ will provide important information about the modification of nuclear parton distributions. We desire that our theoretical results relevant to the NA3 experiment ($E_{beam} = 150$ GeV) can provide useful reference for the forthcoming E906 Sea Quest experiment at Fermi Lab, and give valuable insight in the energy loss of an incoming quark in cold nuclear matter.

**Acknowledgments** This work was supported in part by the National Natural Science Foundation of China(11405043, 11575052).

[1] K. Adcox etal., Phys.Rev.Lett. 88, 022301(2001).

[2] S. S. Adler etal., Phys.Rev.Lett. 91, 072301(2003).

[3] J. Admas etal., Phys.Rev.Lett. 91, 172302(2003).

[4] S. S. Adler etal., Phys.Rev.Lett. 96, 201301(2006).

[5] A. Adare etal., Phys.Rev.Lett. 98, 172301(2007).

[6] A. Adare etal., Phys.Rev.Lett. 104, 252301(2010).

[7] K. Aamodt etal., Phys. Lett. B696, 30(2011).

[8] S. Drell, T. M. Yan, Phys.Rev.Lett. 25, 316(1970).

[9] J. Badier et al., Phys. Lett. B104, 335(1981).

[10] P. Bordal et al., Phys. Lett. B193, 368(1987).

[11] D. M. Alde et al., Phys.Rev.Lett. 64, 2479(1990).

[12] M. A. Vasiliev et al., Phys.Rev.Lett. 83, 2304(1999).

[13] C. G. Duan et al., Eur.Phys.J. C50, 585(2007).

[14] E. Marco, E. Oset, Nucl.Phys. A645, 303(1999).

[15] M. B. Johnson et al., Phys. Rev. C65, 025203(2002).

[16] M. B. Johnson et al., Phys.Rev.Lett. 86, 4483(2001).

[17] F. Arleo, Phys.Lett. B532, 231(2002).

[18] C. T. garvey, J. C. Peng, Phys.Rev.Lett. 90, 090302(2003).

[19] S. J. Brodsky, P. Hoyer, Gavin, Phys. Lett. B298, 165(1993).

[20] C.G. Duan et al., Eur.Phys.J. C29, 557(2003).

[21] C. G. Duan et al., Eur.Phys.J. C39, 179(2005).

[22] C. G. Duan et al., Phys. Rev. C79, 048201(2009).

[23] L. H. Song, C. G. Duan, and N. Liu, Phys. Lett. B708, 68(2012).

[24] M. Hirai, S. Kumano, and M. Miyama, Phys. Rev. D64, 034003(2001).

[25] M. Hirai, S. Kumano, and T. H. Nagai, Phys. Rev. C76, 074028(2004).

[26] D. de Florian, R. Sassot, Phys. Rev. D69, 034003(2001).

[27] K. J. Eskola, H. Paukkunen, and C. A. Salgado, JHEP 0904, 065(2009).

[28] C. A. Salgado, U. A. Wiedemann, Phys. Rev. D68, 014008(2003).

[29] F. Arleo, J. High Energy Phys. 0211, 044(2002).

[30] R. Baier etal., Nucl.Phys. B484, 265(1997).

[31] F. James, CERN Program Library Long Writeup D506.

[32] N. Liu etal., Phys. Lett. B749, 88(2015).

[33] P. E. reimer (Fermilab Sea Quest Collaboration), J. Phys.: Conf. Ser. 295, 0.2011(2011).